



TITLE:

Velocity correlation of a lipid in the lipid-bilayer membrane at the equilibrium

AUTHOR(S):

Inaura, Keiichiro; Fujitani, Youhei

CITATION:

Inaura, Keiichiro ...[et al]. Velocity correlation of a lipid in the lipid-bilayer membrane at the equilibrium. 物性研究 2006, 87(1): 61-62

ISSUE DATE:

2006-10-20

URL:

<http://hdl.handle.net/2433/110652>

RIGHT:

Velocity correlation of a lipid in the lipid-bilayer membrane at the equilibrium

Graduate School of Fundamental Sci & Tech Keio Univ.

Keiichiro Inaura¹, Youhei Fujitani

我々は脂質二重膜上の脂質分子の自己拡散を考える。膜は揺らいでおり、曲げ弾性をもつ圧縮性の2次元流体で、非圧縮 Stokes 流体中に挟まれていると考える。揺動散逸定理から速度場の相関を求め、モード結合理論を用いると、脂質分子の速度相関の表式は速度場を用いて表される。こうして典型的なパラメータに対して得られた速度相関の表式を数値積分した結果によれば、相関関数の時間依存性は時間 t が大きくなるにつれて t^{-1} から $t^{-3/2}$ に遷移することがわかった。

The velocity correlation function (VCF) of a tracer particle decays away as $t^{-d/2}$ as the time t tends to the infinity in a d -dimensional equilibrium fluid[1]. The Brownian particle — a larger impurity particle — also has this long time tail[2]. A biomembrane is a two-dimensional (2D) fluid surrounded with three-dimensional fluids. A membrane protein can be regarded as a Brownian particle[3]. Sera & Rubi[4] showed theoretically that both t^{-1} and $t^{-3/2}$ appear in its velocity correlation[4]. This is reasonable, considering that the momentum on the membrane spread out into the outer fluids. Seki & Komura[5] obtained similar results in a more simple way by introducing a phenomenological momentum relaxation time τ which represents coupling strength between the membrane and the outer fluids. The VCF of the membrane protein was found to decay as $e^{-t/\tau}t^{-3/2}$ or $e^{-t/\tau}t^{-1}$ in the case of the strong- or the weak- coupling limits, respectively.

We study the VCF of the lipid molecule not a membrane protein. We assume that the lipid-bilayer membrane is a compressive 2D Newtonian fluid with the bending rigidity. It fluctuates about the equilibrium in the fluids on its both sides. Using unsteady Stokes approximation, we obtain the Eulerian VCF, $\langle v(\mathbf{k}, t) \cdot v(\mathbf{k}', t') \rangle$, with the aid of the linear response theory[6]. Here, the angular brackets represent the equilibrium ensemble average, while $v(\mathbf{k}, t)$ denotes the Eulerian velocity field with \mathbf{k} representing the wave number vector.

Writing $V(t)$ for a lipid molecule velocity, apart from the fast decaying term, we have

$$\langle V(t) \cdot V(0) \rangle \propto D(t) \equiv \int_0^{k_U} d\mathbf{k} e^{-D_0 k^2 t} \langle v(\mathbf{k}, t) \cdot v(\mathbf{k}, 0) \rangle \quad (1)$$

¹E-mail: nao@beer.appi.keio.ac.jp

in terms of the mode-coupling theory. Here, k_U is a upper cutoff wave number and D_0 is a 'bare' diffusion coefficient, which is independent of the hydrodynamics. We can evaluate its value in terms of the vacancy assisted diffusion[7].

Integrating right-hand side of (1) numerically by use of typical values, we obtain results shown as crosses in Fig.1. Two dotted lines with the slope of -1 and $-\frac{3}{2}$ are shown as guides. We can find that the VCF shifts from t^{-1} to $t^{-\frac{3}{2}}$ as t increases. We are now studying which factor gives the time around which the transition takes place.




Figure 1: The behaviour of $D(t)$

References

- [1] B. J. Alder and T. E. Wainwright, Phys. Rev. A **1** (1970), 18.
- [2] R. Zwanzig and M. Bixon, Phys. Rev. A **2** (1970), 2005.
- [3] P. G. Saffman, J. Fluid Mech. **73** (1967), 593.
- [4] A. Serra and J. M. Rubi, Physica **142A** (1987), 342.
- [5] K. Seki and S. Komura, Phys. Rev. E **47** (1993), 2377.
- [6] Y. Fujitani, J. Chem. Phys. **118** (2002), 7787.
- [7] K. Nakazato and K. Kitahara, Prog. Theor. Phys. **64** (1980), 2261